

- Ulmer, T. S., Ramirez, B. E., Delaglio, F., and Bax, A. (2003). Evaluation of backbone proton positions and dynamics in a small protein by liquid crystal NMR spectroscopy. *J. Am. Chem. Soc.* **125**, 9179–9191.
- Veerapandian, B., Gilliland, G. L., Raag, R., Svensson, A. L., Masui, Y., Hirai, Y., and Poulos, T. L. (1992). Functional implications of interleukin-1-beta based on the 3-dimensional structure. *Proteins* **12**, 10–23.
- Vijay-Kumar, S., Bugg, C. E., and Cook, W. J. (1987). Structure of ubiquitin refined at 1.8 Å resolution. *J. Mol. Biol.* **194**, 531–544.
- Vinson, V. K., Archer, S. J., Lattman, E. E., Pollard, T. D., and Torchia, D. A. (1993). 3-Dimensional solution structure of Acanthamoeba profilin-I. *J. Cell Biol.* **122**, 1277–1283.
- Wand, A. J., Urbauer, J. L., McEvoy, R. P., and Bieber, R. J. (1996). Internal dynamics of human ubiquitin revealed by C-13-relaxation studies of randomly fractionally labeled protein. *Biochemistry* **35**, 6116–6125.
- Weichsel, A., Gasdaska, J. R., Powis, G., and Montfort, W. R. (1996). Crystal structures of reduced, oxidized, and mutated human thioredoxins: Evidence for a regulatory homodimer. *Structure* **4**, 735–751.
- Wu, C. H., Ramamoorthy, A., Gierasch, L. M., and Opella, S. J. (1995). Simultaneous characterization of the amide H-1 chemical shift, H-1-N-15 dipolar, and N-15 chemical-shift interaction tensors in a peptide-bond by 3-dimensional solid-state NMR-spectroscopy. *J. Am. Chem. Soc.* **117**, 6148–6149.
- Zweckstetter, M., and Bax, A. (2000). Prediction of sterically induced alignment in a dilute liquid crystalline phase: Aid to protein structure determination by NMR. *J. Am. Chem. Soc.* **122**, 3791–3792.
- Zweckstetter, M., and Bax, A. (2001). Single-step determination of protein substructures using dipolar couplings: Aid to structural genomics. *J. Am. Chem. Soc.* **123**, 9490–9491.
- Zweckstetter, M., and Bax, A. (2002). Evaluation of uncertainty in alignment tensors obtained from dipolar couplings. *J. Biomol. NMR* **23**, 127–137.
- Zweckstetter, M., Hummer, G., and Bax, A. (2004). Prediction of charge-induced molecular alignment of biomolecules dissolved in dilute liquid-crystalline phases. *Biophys. J.* **86**, 3444–3460.

[4] Rapid NMR Data Collection

By HANUDATTA S. ATREYA and THOMAS SZYPERSKI

Abstract

Rapid data collection is an area of intense research in biomolecular NMR spectroscopy, in particular for high-throughput structure determination in structural genomics. NMR data acquisition and processing protocols for rapidly obtaining high-dimensional spectral information aim at avoiding sampling limited data collection and are reviewed here with emphasis on G-matrix Fourier transform NMR spectroscopy.