

FIRST QUALIFYING EXAM  
August 24, 2000

*There are two sections, Algebra and Analysis. Solve **FOUR** problems from each section. Do **NOT** hand in more than four problems for a section; if you do, only the first four problems will be counted. Do each problem on a separate sheet. Show all work. Be sure to write your person number clearly on **EACH** sheet. Remember, your goal is to convince the grader that you know what you are doing.*

**Section I. Algebra. Do any four problems.**

1. Show that a finite integral domain is a field.
  
2. Consider the conjecture: If a field  $F$  is infinite, then every nontrivial subfield of  $F$  is infinite. Prove or give a counterexample with proof.
  
3. Let  $\varphi: \mathbf{Z} \oplus \mathbf{Z} \rightarrow \mathbf{Z}$  be the group homomorphism defined by  $\varphi(m, n) = 4m - 7n$ . Let  $H$  be the subgroup of  $Z$  generated by the element 5. Find two elements that generate  $\varphi^{-1}(H)$ . (Part of this problem is to prove that the elements actually generate  $\varphi^{-1}(H)$ .)

4. Prove Eisenstein's Criteria. That is, show that if

$$P(x) = a_0 + a_1x + \cdots + a_nx^n$$

is a polynomial over the ring  $\mathbf{Z}$  of integers, and if  $p$  is a prime such that

$$p \text{ divides } a_i, \text{ for } 0 \leq i < n$$

$$p \text{ does not divide } a_n$$

$$p^2 \text{ does not divide } a_0$$

then  $P(x)$  is irreducible over  $\mathbf{Z}$ .

5. Let  $V$  be the vector space of all  $2 \times 2$  matrices with real coefficients. Define an inner product  $\langle \cdot, \cdot \rangle$  on  $V$  by

$$\langle A, B \rangle = \text{trace}(AB^t).$$

Find an orthonormal basis for the orthogonal complement in  $V$  of the  $2 \times 2$  identity matrix.

6. Let  $V$  and  $W$  be finite dimensional vector spaces over  $\mathbf{R}$ . Let  $M$  be a non-zero subspace of  $V$ , and let  $N$  be a subspace of  $W$ . If  $T(M) \subset N$  for all linear transformations  $T: V \rightarrow W$ , prove that  $N = W$ .

7. Let  $M$  be the vector space of upper triangular  $2 \times 2$  matrices, with basis

$$E_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, E_2 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \text{ and } E_3 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}.$$

Let  $T$  be the linear transformation on  $M$  defined by

$$TX = \begin{pmatrix} 1 & 1 \\ 0 & 2 \end{pmatrix} X \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}.$$

Find the matrix  $A = (a_{ij})$  of  $T$ . That is, find the matrix  $A$  that satisfies  $TE_j = \sum_{i=1}^3 a_{ij} E_i$ .

8. A square matrix  $N$  is nilpotent if there is a positive integer  $m$  such that  $N^m = 0$ .

(a) Find a scalar  $\lambda$  and a nilpotent matrix  $N$  such that

$$\begin{pmatrix} 3 & 1 \\ -1 & 1 \end{pmatrix} = \lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + N.$$

(b) Let  $P(x)$  be a polynomial with real coefficients. Use part a) to show that

$$P \left( \begin{pmatrix} 3 & 1 \\ -1 & 1 \end{pmatrix} \right) = \begin{pmatrix} P(2) + P'(2) & P'(2) \\ -P'(2) & P(2) - P'(2) \end{pmatrix}.$$

## Section II. Analysis. Do any four problems.

9. Let  $\sum_{n=0}^{\infty} a_n(x-a)^n$  be an infinite series with radius of convergence  $R > 0$ . Show that if  $0 < r < R$ , then the series converges uniformly for  $|x-a| \leq r$ .

10. Let  $f: [0, 1] \rightarrow \mathbf{R}$  be continuous. Show that

$$\lim_{j \rightarrow \infty} \int_0^1 \cos \left( 2\pi x + \frac{f(x)}{j} \right) dx = 0.$$

11. A set  $X$  in  $\mathbf{R}^n$  is connected if there do not exist open sets  $A$  and  $B$  in  $\mathbf{R}^n$  such that all the following hold:

- (1)  $X \cap A \neq \emptyset$
- (2)  $X \cap B \neq \emptyset$
- (3)  $X \subset (A \cup B)$
- (4)  $X \cap A \cap B = \emptyset$ .

Let  $X$  and  $Y$  be connected subsets of  $\mathbf{R}^N$  such that  $X \cap Y \neq \emptyset$ . Use the definition above to show that  $X \cup Y$  is connected.

12. For each continuously differentiable simple closed curve  $\mathcal{C}$  (oriented counter-clockwise) in  $\mathbf{R}^2$ , define  $\tau(\mathcal{C})$  by

$$\tau(\mathcal{C}) = \int_{\mathcal{C}} 2y^3 dx + (3x - 2x^3) dy.$$

Use Green's Theorem to find the continuously differentiable simple closed curve  $\mathcal{C}$  for which  $\tau(\mathcal{C})$  is maximal.

13. Let  $f: \mathbf{R}^3 \rightarrow \mathbf{R}$  be a  $C^1$  function such that  $f(0, 0, 0) = 0$ . Let  $a, b$ , and  $c$  be real constants such that the derivative of  $f$  at  $(0, 0, 0)$  is  $(a, b, c)$ .

i. Give the minimal conditions on  $a, b$ , and  $c$  that ensure there exists a neighborhood  $N$  of  $(0, 0)$  in  $\mathbf{R}^2$  and a  $C^1$  function  $g: N \rightarrow \mathbf{R}$  such that  $g(0, 0) = 0$  and  $f(x, y, g(x, y)) = 0$  for all  $(x, y)$  in  $N$ .

ii. Assume that suitable conditions on  $a, b$ , and  $c$  are given for part i. Find the derivative of  $g$  at  $(0, 0)$ .

14. Let  $S$  be a compact subset of  $\mathbf{R}^n$ . Prove  $S$  is closed and bounded.