

11 | Tietze Extension Theorem

11.1 Tietze Extension Theorem (v.1). *Let X be a normal space, let $A \subseteq X$ be a closed subspace, and let $f: A \rightarrow [a, b]$ be a continuous function for some $[a, b] \subseteq \mathbb{R}$. There exists a continuous function $\tilde{f}: X \rightarrow [a, b]$ such that $\tilde{f}|_A = f$.*

11.2 Definition. Let X, Y be topological spaces and let $\{f_n: X \rightarrow Y\}$ be a sequence of functions. We say that the sequence $\{f_n\}$ *converges pointwise* to a function $f: X \rightarrow Y$ if for each $x \in X$ the sequence $\{f_n(x)\} \subseteq Y$ converges to the point $f(x)$.

11.4 Definition. Let X be a topological space, let (Y, ϱ) be a metric space, and let $\{f_n: X \rightarrow Y\}$ be a sequence of functions. We say that the sequence $\{f_n\}$ *converges uniformly* to a function $f: X \rightarrow Y$ if for every $\varepsilon > 0$ there exists $N > 0$ such that

$$\varrho(f(x), f_n(x)) < \varepsilon$$

for all $x \in X$ and for all $n > N$.

11.5 Note. If a sequence $\{f_n\}$ converges uniformly to f then it also converges pointwise to f , but the converse is not true in general.

11.6 Proposition. *Let X be a topological space and let (Y, ϱ) be a metric space. Assume that $\{f_n: X \rightarrow Y\}$ is a sequence of functions that converges uniformly to $f: X \rightarrow Y$. If all functions f_n are continuous then f is also a continuous function.*

11.7 Lemma. *Let X be a normal space, $A \subseteq X$ be a closed subspace, and let $f: A \rightarrow \mathbb{R}$ be a continuous function such that for some $C > 0$ we have $|f(x)| \leq C$ for all $x \in A$. There exists a continuous function $g: X \rightarrow \mathbb{R}$ such that $|g(x)| \leq \frac{1}{3}C$ for all $x \in X$ and $|f(x) - g(x)| \leq \frac{2}{3}C$ for all $x \in A$.*

11.8 Tietze Extension Theorem (v.2). *Let X be a normal space, let $A \subseteq X$ be a closed subspace, and let $f: A \rightarrow \mathbb{R}$ be a continuous function. There exists a continuous function $\tilde{f}: X \rightarrow \mathbb{R}$ such that $\tilde{f}|_A = f$.*

11.9 Theorem. *Let X be a space satisfying T_1 . The following conditions are equivalent:*

- 1) *X is a normal space.*
- 2) *For any closed sets $A, B \subseteq X$ such that $A \cap B = \emptyset$ there is a continuous function $f: X \rightarrow [0, 1]$ such that $A \subseteq f^{-1}(\{0\})$ and $B \subseteq f^{-1}(\{1\})$.*
- 3) *If $A \subseteq X$ is a closed set then any continuous function $f: A \rightarrow \mathbb{R}$ can be extended to a continuous function $\tilde{f}: X \rightarrow \mathbb{R}$.*