

11 | Tietze Extension Theorem

The main goal of this chapter is to prove the following fact which describes one of the most useful properties of normal spaces:

11.1 Tietze Extension Theorem (v.1). *Let X be a normal space, let $A \subseteq X$ be a closed subspace, and let $f: A \rightarrow [a, b]$ be a continuous function for some $[a, b] \subseteq \mathbb{R}$. There exists a continuous function $\tilde{f}: X \rightarrow [a, b]$ such that $\tilde{f}|_A = f$.*

The main idea of the proof is to use Urysohn Lemma 10.1 to construct functions $\tilde{f}_n: X \rightarrow [a, b]$ for $n = 1, 2, \dots$ such that as n increases $\tilde{f}_n|_A$ gives ever closer approximations of f . Then we take \tilde{f} to be the limit of the sequence $\{\tilde{f}_n\}$. We start by looking at sequences of functions and their convergence.

11.2 Definition. Let X, Y be a topological spaces and let $\{f_n: X \rightarrow Y\}$ be a sequence of functions. We say that the sequence $\{f_n\}$ *converges pointwise* to a function $f: X \rightarrow Y$ if for each $x \in X$ the sequence $\{f_n(x)\} \subseteq Y$ converges to the point $f(x)$.

11.3 Note. If $\{f_n: X \rightarrow Y\}$ is a sequence of continuous functions that converges pointwise to $f: X \rightarrow Y$ then f need not be continuous. For example, let $f_n: [0, 1] \rightarrow \mathbb{R}$ be the function given by $f_n(x) = x^n$. Notice that $f_n(x) \rightarrow 0$ for all $x \in [0, 1)$ and that $f_n(1) \rightarrow 1$. Thus the sequence $\{f_n\}$ converges pointwise to the function $f: [0, 1] \rightarrow \mathbb{R}$ defined by

$$f(x) = \begin{cases} 0 & \text{for } x \neq 1 \\ 1 & \text{for } x = 1 \end{cases}$$

The functions f_n are continuous but f is not.

11.4 Definition. Let X be a topological space, let (Y, ρ) be a metric space, and let $\{f_n: X \rightarrow Y\}$ be a sequence of functions. We say that the sequence $\{f_n\}$ *converges uniformly* to a function $f: X \rightarrow Y$ if

for every $\varepsilon > 0$ there exists $N > 0$ such that

$$\varrho(f(x), f_n(x)) < \varepsilon$$

for all $x \in X$ and for all $n > N$.


11.5 Note. If a sequence $\{f_n\}$ converges uniformly to f then it also converges pointwise to f , but the converse is not true in general.

11.6 Proposition. Let X be a topological space and let (Y, ϱ) be a metric space. Assume that $\{f_n: X \rightarrow Y\}$ is a sequence of functions that converges uniformly to $f: X \rightarrow Y$. If all functions f_n are continuous then f is also a continuous function.

Proof. Let $U \subseteq Y$ be an open set. We need to show that the set $f^{-1}(U) \subseteq X$ is open. It suffices to check that each point $x_0 \in f^{-1}(U)$ has an open neighborhood V such that $V \subseteq f^{-1}(U)$. Since U is an open set there exists $\varepsilon > 0$ such $B(f(x_0), \varepsilon) \subseteq U$. Choose $N > 0$ such that $\varrho(f(x), f_N(x)) < \frac{\varepsilon}{3}$ for all $x \in X$, and take $V = f_N^{-1}(B(f_N(x_0), \frac{\varepsilon}{3}))$. Since f_N is a continuous function the set V is an open neighborhood of x_0 in X . It remains to show that $V \subseteq f^{-1}(U)$. For $x \in V$ we have:

$$\varrho(f(x), f(x_0)) \leq \varrho(f(x), f_N(x)) + \varrho(f_N(x), f_N(x_0)) + \varrho(f_N(x_0), f(x_0)) < \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} = \varepsilon$$

This means that $f(x) \in B(f(x_0), \varepsilon) \subseteq U$, and so $x \in f^{-1}(U)$.



□

11.7 Lemma. Let X be a normal space, $A \subseteq X$ be a closed subspace, and let $f: A \rightarrow \mathbb{R}$ be a continuous function such that for some $C > 0$ we have $|f(x)| \leq C$ for all $x \in A$. There exists a continuous function $g: X \rightarrow \mathbb{R}$ such that $|g(x)| \leq \frac{1}{3}C$ for all $x \in X$ and $|f(x) - g(x)| \leq \frac{2}{3}C$ for all $x \in A$.

Proof. Define $Y := f^{-1}([-C, -\frac{1}{3}C])$, $Z := f^{-1}([\frac{1}{3}C, C])$. Since $f: A \rightarrow \mathbb{R}$ is a continuous function these sets are closed in A , but since A is closed in X the sets Y and Z are also closed in X . Since $Y \cap Z = \emptyset$ by the Urysohn Lemma 10.1 there is a continuous function $h: X \rightarrow [0, 1]$ such that $h(Y) \subseteq \{0\}$ and $h(Z) \subseteq \{1\}$. Define $g: X \rightarrow \mathbb{R}$ by

$$g(x) := \frac{2C}{3} \left(h(x) - \frac{1}{2} \right)$$

□

Proof of Theorem 11.1. Without loss of generality we can assume that $[a, b] = [0, 1]$. For $n = 1, 2, \dots$ we will construct continuous functions $g_n: X \rightarrow \mathbb{R}$ such that

- (i) $|g_n(x)| \leq \frac{1}{3} \cdot \left(\frac{2}{3}\right)^{n-1}$ for all $x \in X$;
- (ii) $\left|f(x) - \sum_{i=1}^n g_i(x)\right| \leq \left(\frac{2}{3}\right)^n$ for all $x \in A$.

We argue by induction. Existence of g_1 follows directly from Lemma 11.7. Assume that for some $n \geq 1$ we already have functions g_1, \dots, g_n satisfying (i) and (ii). In Lemma 11.7 take f to be the function $f - \sum_{i=1}^n g_i$ and take $C = \left(\frac{2}{3}\right)^n$. Then we can take $g_{n+1} := g$ where g is the function given by the lemma.

Let $\tilde{f}_n := \sum_{i=1}^n g_i$ and let $\tilde{f} := \sum_{i=1}^{\infty} g_i$. Using condition (i) we obtain that the sequence $\{\tilde{f}_n\}$ converges uniformly to \tilde{f} (exercise). Since each of the functions \tilde{f}_n is continuous, thus by Proposition 11.6 we obtain that \tilde{f} is a continuous function. Also, using (ii) we obtain that $\tilde{f}(x) = f(x)$ for all $x \in A$ (exercise). □

Here is another useful reformulation of Tietze Extension Theorem:

11.8 Tietze Extension Theorem (v.2). *Let X be a normal space, let $A \subseteq X$ be a closed subspace, and let $f: A \rightarrow \mathbb{R}$ be a continuous function. There exists a continuous function $\tilde{f}: X \rightarrow \mathbb{R}$ such that $\tilde{f}|_A = f$.*

Proof. It is enough to show that for any continuous function $g: A \rightarrow (-1, 1)$ we can find a continuous function $\tilde{g}: X \rightarrow (-1, 1)$ such that $\tilde{g}|_A = g$. Indeed, if this holds then given a function $f: A \rightarrow \mathbb{R}$ let $g = hf$ where $h: \mathbb{R} \rightarrow (-1, 1)$ is an arbitrary homeomorphism. Then we can take $\tilde{f} = h^{-1}\tilde{g}$.

Assume then that $g: A \rightarrow (-1, 1)$ is a continuous function. By Theorem 11.1 there is a function $g_1: X \rightarrow [-1, 1]$ such that $g_1|_A = g$. Let $B := g_1^{-1}(\{-1, 1\})$. The set B is closed in X and $A \cap B = \emptyset$ since $g_1(A) = g(A) \subseteq (-1, 1)$. By Urysohn Lemma 10.1 there is a continuous function $k: X \rightarrow [0, 1]$ such that $B \subseteq k^{-1}(\{0\})$ and $A \subseteq k^{-1}(\{1\})$. Let $\tilde{g}(x) := k(x) \cdot g_1(x)$. We have:

- 1) if $g_1(x) \in (-1, 1)$ then $\tilde{g}(x) \in (-1, 1)$
- 2) if $g_1(x) \in \{-1, 1\}$ then $x \in B$ so $\tilde{g}(x) = 0 \cdot g_1(x) = 0$

It follows that $\tilde{g}: X \rightarrow (-1, 1)$. Also, \tilde{g} is a continuous function since k and g_1 are continuous. Finally, if $x \in A$ then $\tilde{g}(x) = 1 \cdot g_1(x) = g(x)$, so $\tilde{g}|_A = g$. □

Tietze Extension Theorem holds for functions defined on normal spaces. It turns out the function extension property is actually equivalent to the notion of normality of a space:

11.9 Theorem. *Let X be a space satisfying T_1 . The following conditions are equivalent:*

- 1) X is a normal space.

- 2) For any closed sets $A, B \subseteq X$ such that $A \cap B = \emptyset$ there is a continuous function $f: X \rightarrow [0, 1]$ such that $A \subseteq f^{-1}(\{0\})$ and $B \subseteq f^{-1}(\{1\})$.
- 3) If $A \subseteq X$ is a closed set then any continuous function $f: A \rightarrow \mathbb{R}$ can be extended to a continuous function $\tilde{f}: X \rightarrow \mathbb{R}$.

Proof. The implication 1) \Rightarrow 2) is the Urysohn Lemma 10.1 and 2) \Rightarrow 1) is Proposition 9.15. The implication 1) \Rightarrow 3) is the Tietze Extension Theorem 11.8. The proof of implication 3) \Rightarrow 1) is an exercise.

□

Exercises to Chapter 11

E11.1 Exercise. Prove implication 3) \Rightarrow 1) of Theorem 11.9.

E11.2 Exercise. Let X be a normal space, let $A \subseteq X$ be a closed subspace, and let $f: A \rightarrow \mathbb{R}$ be a continuous function.

- a) Assume that $g: X \rightarrow \mathbb{R}$ is a continuous function such that $f(x) \leq g(x)$ for all $x \in A$. Show that there exists a continuous function $F: X \rightarrow \mathbb{R}$ satisfying $F|_A = f$ and $F(x) \leq g(x)$ for all $x \in X$.
- b) Assume that $g, h: X \rightarrow \mathbb{R}$ are a continuous function such that $h(x) \leq f(x) \leq g(x)$ for all $x \in A$ and $h(x) \leq g(x)$ for all $x \in X$. Show that there exists a continuous function $F': X \rightarrow \mathbb{R}$ satisfying $F'|_A = f$ and $h(x) \leq F'(x) \leq g(x)$ for all $x \in X$.

E11.3 Exercise. Recall that if X is a topological space then a subspace $Y \subseteq X$ is called a retract of X if there exists a continuous function $r: X \rightarrow Y$ such that $r(x) = x$ for all $x \in Y$. Let X be a normal space and let $Y \subseteq X$ be a closed subspace of X such that $Y \cong \mathbb{R}$. Show that Y is a retract of X .

E11.4 Exercise. Let X be topological space. Recall from Exercise 10.3 that a set $A \subseteq X$ is a G_δ -set if there exists a countable family of open sets U_1, U_2, \dots such that $A = \bigcap_{n=1}^{\infty} U_n$.

- a) Show that if X is a normal space and $A \subseteq X$ is a closed G_δ -set then there exists a continuous function $f: X \rightarrow [0, 1]$ such that $A = f^{-1}(\{0\})$.
- b) Show that if X is a normal space and $A, B \subseteq X$ are closed G_δ -sets such that $A \cap B = \emptyset$ then there exists a continuous function $f: X \rightarrow [0, 1]$ such that $A = f^{-1}(\{0\})$ and $B = f^{-1}(\{1\})$.